

Medical Policy:

Colony Stimulating Factors: Fulphila™ (pegfilgrastim-jmdb)

POLICY NUMBER	LAST REVIEW	ORIGIN DATE
MG.MM.PH.61	March 21, 2024	

Medical Guideline Disclaimer Property of EmblemHealth. All rights reserved.

The treating physician or primary care provider must submit to EmblemHealth, or ConnectiCare, as applicable (hereinafter jointly referred to as “EmblemHealth”), the clinical evidence that the member meets the criteria for the treatment or surgical procedure. Without this documentation and information, EmblemHealth will not be able to properly review the request preauthorization or post-payment review. The clinical review criteria expressed below reflects how EmblemHealth determines whether certain services or supplies are medically necessary. This clinical policy is not intended to pre-empt the judgment of the reviewing medical director or dictate to health care providers how to practice medicine. Health care providers are expected to exercise their medical judgment in rendering appropriate care. Health care providers are expected to exercise their medical judgment in rendering appropriate care.

EmblemHealth established the clinical review criteria based upon a review of currently available clinical information (including clinical outcome studies in the peer reviewed published medical literature, regulatory status of the technology, evidence-based guidelines of public health and health research agencies, evidence-based guidelines and positions of leading national health professional organizations, views of physicians practicing in relevant clinical areas, and other relevant factors). EmblemHealth expressly reserves the right to revise these conclusions as clinical information changes and welcomes further relevant information. Each benefit program defines which services are covered. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered and/or paid for by EmblemHealth, as some programs exclude coverage for services or supplies that EmblemHealth considers medically necessary.

If there is a discrepancy between this guideline and a member's benefits program, the benefits program will govern. Identification of selected brand names of devices, tests and procedures in a medical coverage policy is for reference only and is not an endorsement of any one device, test or procedure over another. In addition, coverage may be mandated by applicable legal requirements of a state, the Federal Government or the Centers for Medicare & Medicaid Services (CMS) for Medicare and Medicaid members. All coding and web site links are accurate at time of publication.

EmblemHealth may also use tools developed by third parties, such as the MCG™ Care Guidelines, to assist us in administering health benefits. The MCG™ Care Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice. EmblemHealth Services Company, LLC, has adopted this policy in providing management, administrative and other services to EmblemHealth Plan, Inc., EmblemHealth Insurance Company, EmblemHealth Services Company, LLC, and Health Insurance Plan of Greater New York (HIP) related to health benefit plans offered by these entities. ConnectiCare, an EmblemHealth company, has also adopted this policy. All of the aforementioned entities are affiliated companies under common control of EmblemHealth Inc.

Definitions

Fulphila is a colony stimulating factor that acts on hematopoietic cells by binding to specific cell surface receptors thereby stimulating proliferation, differentiation, commitment, and end cell functional activation

Length of Authorization

Coverage will be provided for six months and may be renewed.

Dosing Limits [Medical Benefit]

Max Units (per dose and over time)

- Acute radiation exposure
 - 12 billable units weekly for 2 doses
- All other indications:
 - 12 billable units per 14 days for all other indications

Guideline

I. Initial Approval Criteria

Fulphila may be considered medically necessary if one of the below conditions are met **AND** use is consistent with the medical necessity criteria that follows:

Neulasta and Udenyca are the preferred agents for Commercial, Medicaid, and Medicare members.

1. The patient has failed treatment with Neulasta **AND** Udenyca or they are contraindicated ††; **OR**
2. The patient is continuing previously established therapy with Fulphila for their current chemotherapy regimen; **AND**
3. A member does not have access to, or benefits for, home health services; **OR**
4. A member is expected to receive G-CSF for 5 consecutive days or more; **OR**
5. Fulphila is used in combination with one of the following chemotherapy regimens*:
 - A. Bladder Cancer:
 - i. Dose dense MVAC (methotrexate, vinblastine, doxorubicin, cisplatin)
 - B. Breast Cancer:
 - i. Dose dense AC followed by T (doxorubicin, cyclophosphamide, paclitaxel)
 - C. Non-Hodgkin's Lymphoma:
 - i. Dose dense CHOP-14 (cyclophosphamide, doxorubicin, vincristine, prednisone)

†† Commercial, Medicaid, AND Medicare members are subject to this step therapy

* Pegylated filgrastim is the only G-CSF product used in the published clinical trials for these regimens. The requesting provider should provide journal citations supporting this request for regimens other than those listed.

Coverage for Fulphila™ (pegfilgrastim-jmdb) is provided in the following conditions:

Prophylactic use in patients with non-myeloid malignancy†

1. Patient is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia of 20% or greater § ; **OR**
2. Patient is undergoing myelosuppressive chemotherapy with an expected incidence of febrile neutropenia of 10% or greater § **AND** one or more of the following co-morbidities:
 - a. Elderly patients (age 65 or older)
 - b. History of recurrent febrile neutropenia from chemotherapy
 - c. Extensive prior exposure to chemotherapy
 - d. Previous exposure of pelvis, or other areas of large amounts of bone marrow, to radiation
 - e. Pre-existing neutropenia (ANC ≤ 1000/mm³) or bone marrow involvement with tumor
 - f. Patient has a condition that can potentially increase the risk of serious infection (i.e. HIV/AIDS)
 - g. Infection/open wounds
 - h. Recent surgery
 - i. Poor performance status
 - j. Poor renal function (creatinine clearance <50)
 - k. Liver dysfunction (elevated bilirubin >2.0)
 - l. Chronic immunosuppression in the post-transplant setting including organ transplant

Patient who experienced a neutropenic complication from a prior cycle of the same chemotherapy ‡

Patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome) ‡

†FDA-labeled indication, ‡ Compendia recommended indication

§ expected incidence of febrile neutropenia percentages for myelosuppressive chemotherapy regimens can be found in the NCCN Myeloid Growth Factors Clinical Practice Guideline at NCCN.org

II. Renewal Criteria

Same as initial prior authorization policy criteria

III. Dosage/Administration

Indication	Dose
Acute radiation exposure	6 mg subcutaneously weekly for 2 doses (Use weight based dosing for pediatrics weight < 45 kg)
All other indications	6 mg subcutaneously once per chemotherapy cycle and dosed no more frequently than every 14 days (Use weight based dosing for pediatric patients weighing less than 45 kg)

*Do not administer within 14 days before and 24 hours after administration of cytotoxic chemotherapy

Applicable Procedure Codes

Code	Description
Q5108	Injection, pegfilgrastim-jmdb, biosimilar, (Fulphila), 0.5 mg

Applicable NDCs

Code	Description
67457-0833-06	Fulphila 6 mg prefilled syringe

ICD-10 Diagnoses

Code	Description
D70.1	Agranulocytosis secondary to cancer chemotherapy
D70.9	Neutropenia, unspecified
T45.1X5A	Adverse effect of antineoplastic and immunosuppressive drugs initial encounter
T45.1X5D	Adverse effect of antineoplastic and immunosuppressive drugs subsequent encounter
T45.1X5S	Adverse effect of antineoplastic and immunosuppressive drugs sequela
Z41.8	Encounter for other procedures for purposes other than remedying health state
Z48.290	Encounter for aftercare following bone marrow transplant
Z51.11	Encounter for antineoplastic chemotherapy

Z51.12	Encounter for antineoplastic immunotherapy
Z51.89	Encounter for other specified aftercare
Z52.001	Unspecified donor, stem cells
Z52.011	Autologous donor, stem cells
ZZ52.091	Other blood donor, stem cells
Z94.81	Bone marrow transplant status
Z94.84	Stem cells transplant status

Revision History

Company(ies)	DATE	REVISION
EmblemHealth & ConnectiCare	3/21/2024	Annual Review: updated dosing chart
EmblemHealth & ConnectiCare	9/13/2023	Annual Review: no criteria changes
EmblemHealth & ConnectiCare	4/07/2022	Transferred policy to new template
EmblemHealth & ConnectiCare	1/1/2021	Extended coverage duration from 4 to 6 months
EmblemHealth & ConnectiCare	11/2/2020	Effective 01/01/2021 , Member must fail trial of Neulasta AND Udenyca, prior to using Fulphila (Medicare members are subject to this step therapy).
EmblemHealth & ConnectiCare	11/20/2019	Neulasta and Udenyca are the preferred agents for Medicare members. (Step protocol not mandated for Medicare members).
EmblemHealth & ConnectiCare	12/18/2018	Added Step therapy to use Neulasta AND Udenyca prior to initiating Fulphila therapy.

References

1. Fulphila [package insert]. Mylan GmbH, Zurich, Switzerland. May 2019